


## aktiv in ganz Europa

## LS Risk Score Assessment

Reasons for a biopsy: Suspicion of a precancer Uncertain diagnosis



| Medical Assessment: Clinical Score |      |      |            |
|------------------------------------|------|------|------------|
|                                    | None | Some | Widespread |
| Erosions                           | 0    | 1    | 2          |
| Hyperkeratosis                     | 0    | 1    | 2          |
| Rhagades                           | 0    | 1    | 2          |
| Synechiae                          | 0    |      | 2          |
| Stenosis                           | 0    | 1    | 2          |
| Atrophy                            | 0    | 1    | 2          |

If the LS score is high but there is no suspicion of a precancerous stage, the biopsy can be dispensed with.

A score of 4: the probability of LS >90%



Günthert et al. 2012 *J Sex Med* Naswa et al. 2015 *Ind J Sex Transm Dis* 

www.gyn-zentrum.ch

## Glossary:

Erosions: Non-bleeding loss of skin or mucous membrane.

Hyperkeratosis: Excessive cornification of large or small areas of the outermost layer of the epidermis.

Rhagades: Slit-like lacerations on the skin, that heal without scars, and that result from overstretching

due to reduced elasticity (e.g. due to dehydration, cold or neurodermatitis).

Synechiae: The adhesion of two tissue layers in the form of connective tissue, that are normally

separated from each other.

Stenosis: Constriction.

Atrophy: Decrease in volume or size.

Please also see the following tabs on our website www.lichensclerosus.ch

"Biopsies - necessary / appropriate?"

"Facts about cortisone"

Verein Lichen Sclerosus, Bleicheweg 6, CH - 5605 - 5605 Dottikon, www.lichensclerosus.ch, www.juckenundbrennen.eu